

- (1) Consider  $A = \{q \in \mathbb{Q} : q^2 \geq 2\}$  as a subset of the metric space  $(\mathbb{Q}, d)$ , where  $d(x, y) = |x - y|$ . Then  $A$  is
- A) closed but not open in  $\mathbb{Q}$
  - B) open but not closed in  $\mathbb{Q}$
  - C) neither open nor closed in  $\mathbb{Q}$
  - D) both open and closed in  $\mathbb{Q}$ .
- (2) The set  $\mathbb{N}$  considered as a subspace of  $(\mathbb{R}, d)$  where  $d(x, y) = |x - y|$ , is
- A) closed but not complete
  - B) complete but not closed
  - C) both closed and complete
  - D) neither closed nor complete.
- (3) Let  $Y$  be a totally bounded subset of a metric space  $X$ . Then the closure  $\overline{Y}$  of  $Y$
- A) is totally bounded
  - B) may not be totally bounded even if  $X$  is complete
  - C) is totally bounded if and only if  $X$  is complete
  - D) is totally bounded if and only if  $X$  is compact.
- (4) Let  $X, Y$  be metric spaces,  $f : X \rightarrow Y$  be a continuous function,  $A$  be a bounded subset of  $X$  and let  $B = f(A)$ . Then  $B$  is
- A) bounded
  - B) bounded if  $A$  is also closed
  - C) bounded if  $A$  is compact
  - D) bounded if  $A$  is complete.
- (5) Let  $X$  be a connected metric space and  $U$  be an open subset of  $X$ . Then
- A)  $U$  cannot be closed in  $X$
  - B) if  $U$  is closed in  $X$ , then  $U = X$
  - C) if  $U$  is closed in  $X$ , then  $U = \phi$ , the empty set
  - D) if  $U$  is closed in  $X$  and  $U$  is non-empty, then  $U = X$ .
- (6) Let  $X$  be a connected metric space and  $f : X \rightarrow \mathbb{R}$  be a continuous function. Then  $f(X)$
- A) is whole of  $\mathbb{R}$
  - B) is a bounded subset of  $\mathbb{R}$
  - C) is an interval in  $\mathbb{R}$
  - D) may not be an interval in  $\mathbb{R}$ .

(7) Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined as

$$f(x, y) = \begin{cases} \frac{xy^2}{x^2+y^4}, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases}$$

Let  $D_u f(0, 0)$  denote the directional derivative of  $f$  at  $(0, 0)$  in the direction  $u = (u_1, u_2) \neq (0, 0)$ . Then  $f$  is

- A) continuous at  $(0, 0)$  and  $D_u f(0, 0)$  exist for all  $u$
- B) continuous at  $(0, 0)$  but  $D_u f(0, 0)$  does not exist for some  $u \neq (0, 0)$
- C) not continuous at  $(0, 0)$  but  $D_u f(0, 0)$  exist for all  $u$
- D) not continuous at  $(0, 0)$  and  $D_u f(0, 0)$  does not exist for some  $u \neq (0, 0)$ .

(8) Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be defined as

$$f(x, y) = \frac{x^2 - y^2}{1 + x^2 + y^2}$$

Then

- A)  $\frac{\partial^2 f}{\partial x \partial y}(0, 0)$  and  $\frac{\partial^2 f}{\partial y \partial x}(0, 0)$  exist but are not equal
- B)  $\frac{\partial^2 f}{\partial x \partial y}(0, 0)$  exist but  $\frac{\partial^2 f}{\partial y \partial x}(0, 0)$  does not exist
- C)  $\frac{\partial^2 f}{\partial y \partial x}(0, 0)$  exist but  $\frac{\partial^2 f}{\partial x \partial y}(0, 0)$  does not exist
- D)  $\frac{\partial^2 f}{\partial x \partial y}(0, 0)$  and  $\frac{\partial^2 f}{\partial y \partial x}(0, 0)$  exist and are equal.

(9) The sequence

$$\left\langle \frac{2^{n+1} + 3^{n+1}}{2^n + 3^n} \right\rangle$$

converges to

- A) 1
- B) 2
- C) 3
- D) 5.

(10) The limit of the sequence  $\langle \sqrt{(n+1)(n+2)} - n \rangle$  as  $n \rightarrow \infty$  is

- A)  $\sqrt{2} - 1$
- B) 3
- C)  $3/2$
- D) 0.

(11) The radius of convergence of the series

$$\sum_{n=0}^{\infty} \frac{1}{2^n} x^{3n}$$

is

- A) 1
- B)  $\infty$
- C)  $1/2$
- D)  $2^{1/3}$ .

(12) Which one of the following sequence converges uniformly on the indicated set?

- A)  $f_n(x) = (1 - |x|)^n; \quad x \in (-1, 1)$

- B)  $f_n(x) = \frac{1}{n} \sin nx; \quad x \in \mathbb{R}$   
 C)  $f_n(x) = x^n; \quad x \in [0, 1]$   
 D)  $f_n(x) = \frac{1}{1+x^n}; \quad x \in [0, \infty)$ .

(13) Which one of the following integrals is convergent?

- A)  $\int_1^\infty \frac{1}{x^2} dx$                       B)  $\int_1^\infty \frac{1}{\sqrt{x}} dx$   
 C)  $\int_0^1 \frac{1}{x^2} dx$                       D)  $\int_0^\infty \frac{1}{\sqrt{x}} dx$ .

(14) The value of the integral

$$\int_{-\infty}^{\infty} e^{-x^2} dx$$

is

- A) 0                      B)  $\sqrt{2\pi}$                       C)  $\sqrt{\pi}$                       D)  $\sqrt{\pi/2}$ .

(15) Let  $f : I \rightarrow \mathbb{R}$  be an increasing function where  $I$  is an interval in  $\mathbb{R}$ . Then

- A)  $f^2$  is always increasing  
 B)  $f^2$  is always decreasing  
 C)  $f^2$  is constant  $\Rightarrow f$  is constant  
 D)  $f^2$  may be neither decreasing nor increasing.

(16) Consider the function  $f(x) = x^2$  on  $[0, 1]$  and the partition  $P$  of  $[0, 1]$  given by

$$P = \left\{ 0 < \frac{1}{n} < \frac{2}{n} < \dots < \frac{n-1}{n} < 1 \right\}.$$

Then the upper and the lower Riemann sums of  $f$  are

- A)  $U(f, P) = (1 + \frac{1}{n})(2 - \frac{1}{n})/6$  and  $L(f, P) = (1 + \frac{1}{n})(2 + \frac{1}{n})/6$   
 B)  $U(f, P) = (1 + \frac{1}{n})(2 + \frac{1}{n})/6$  and  $L(f, P) = (1 - \frac{1}{n})(2 - \frac{1}{n})/6$   
 C)  $U(f, P) = (1 + \frac{1}{n})(2 + \frac{1}{n})/6$  and  $L(f, P) = (1 - \frac{1}{n})(2 + \frac{1}{n})/6$   
 D)  $U(f, P) = (1 - \frac{1}{n})(2 + \frac{1}{n})/6$  and  $L(f, P) = (1 + \frac{1}{n})(2 - \frac{1}{n})/6$ .

(17) Which one of the following is true?

- A) If  $\sum a_n$  diverges and  $a_n > 0$ , then  $\sum \frac{a_n}{1+a_n}$  diverges  
 B) If  $\sum a_n$  and  $\sum b_n$  diverge, then  $\sum (a_n + b_n)$  diverges  
 C) If  $\sum a_n$  and  $\sum b_n$  diverge, then  $\sum (a_n + b_n)$  converges  
 D) If  $\sum a_n$  converges and  $\sum b_n$  diverges, then  $\sum (a_n + b_n)$  converges.

(18) If  $\sum a_n = A$ ,  $\sum |a_n| = B$  and  $A$  and  $B$  are finite, then

- A)  $|A| = B$                       B)  $A \leq B$   
 C)  $|A| \geq B$                       D)  $A = B$ .

(19) If  $x_n = 1 + (-1)^n + \frac{1}{2^n}$ , then

- A)  $\limsup x_n = 1$
- B)  $\liminf x_n = 1$
- C)  $x_n$  is a convergent sequence
- D)  $\limsup x_n \neq \liminf x_n$ .

(20) Let  $\langle x_n \rangle$  be the sequence defined by  $x_1 = 2$  and  $x_{n+1} = \frac{1}{2}(x_n + \frac{2}{x_n})$ . Then

- A)  $\langle x_n \rangle$  converges to rational number
- B)  $\langle x_n \rangle$  is an increasing sequence
- C)  $\langle x_n \rangle$  converges to  $2\sqrt{2}$
- D)  $\langle x_n \rangle$  is a decreasing sequence.

(21) Which one of the following series converges?

- A)  $\sum \cos \frac{1}{n^2}$
- B)  $\sum \sin \frac{1}{n^2}$
- C)  $\sum \frac{1}{n^{1+1/n}}$
- D)  $\sum n^{\cos 3}$ .

(22) The sum of the series

$$\sum_{k=1}^{\infty} \frac{1}{(2k-1)^2}$$

is

- A)  $\frac{\pi^2}{8}$
- B)  $\frac{\pi^2}{6}$
- C)  $\frac{\pi}{2}$
- D) 1.

(23) Which one of the following set is not countable?

- A)  $\mathbb{N}^r$ , where  $r \geq 1$  and  $\mathbb{N}$  is the set of natural numbers
- B)  $\{0, 1\}^{\mathbb{N}}$ , the set of all the sequences which takes values 0 and 1
- C)  $\mathbb{Z}$ , set of integers
- D)  $\sqrt{2}\mathbb{Q}$ ,  $\mathbb{Q}$  is set of rational numbers.

(24) Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a continuous function such that  $f(x^2) = f(x)$  for all  $x \in [0, 1]$ . Which one of the following is not true in general?

- A)  $f$  is constant
- B)  $f$  is uniformly continuous
- C)  $f$  is differentiable
- D)  $f(x) \geq 0 \forall x \in [0, 1]$ .

(25) Let  $f : [0, 1] \rightarrow [0, 1]$  be a continuous function and  $I : [0, 1] \rightarrow [0, 1]$  be the identity function. Then  $f$  and  $I$

- A) agree exactly at one point
- B) agree at least at one point
- C) may not agree at any point
- D) agree at most at one point.

(26) For  $x \in \mathbb{R}$ , let  $[x]$  denote the greatest integer  $n$  such that  $n \leq x$ . The function  $h(x) = x[x]$  is

- A) continuous everywhere
- B) continuous only at  $x = \pm 1, \pm 2, \pm 3, \dots$
- C) continuous if  $x \neq \pm 1, \pm 2, \pm 3, \dots$
- D) bounded on  $\mathbb{R}$ .

(27) Let  $\langle x_n \rangle$  be an unbounded sequence in  $\mathbb{R}$ . Then

- A)  $\langle x_n \rangle$  has a convergent subsequence
- B)  $\langle x_n \rangle$  has a subsequence  $\langle x_{n_k} \rangle$  such that  $x_{n_k} \rightarrow 0$
- C)  $\langle x_n \rangle$  has a subsequence  $\langle x_{n_k} \rangle$  such that  $\frac{1}{x_{n_k}} \rightarrow 0$
- D) Every subsequence of  $\langle x_n \rangle$  is unbounded.

(28) Consider the function  $g : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$g(x) = \begin{cases} 0, & \text{if } x \geq 0, \\ e^{-1/x^2}, & \text{if } x < 0. \end{cases}$$

Which one of the following is not true?

- A)  $g$  has derivatives of all orders at every point
- B)  $g^n(0) = 0$  for all  $n \in \mathbb{N}$
- C) Taylor Series expansion of  $g$  about  $x = 0$  converges to  $g$  for all  $x$
- D) Taylor Series expansion of  $g$  about  $x = 0$  converges to  $g$  for all  $x \geq 0$ .

(29) The function

$$f(x) = x \sin x + \frac{1}{1+x^2}; \quad x \in I$$

where  $I \subseteq \mathbb{R}$  is

- A) uniformly continuous if  $I = \mathbb{R}$
- B) uniformly continuous if  $I$  is compact
- C) uniformly continuous if  $I$  is closed
- D) not uniformly continuous on  $[0, 1]$ .

(30) Let the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as

$$f(x) = \begin{cases} x^2, & \text{if } x \in (0, 2) \cap \mathbb{Q}, \\ 2x - 1, & \text{if } x \in (0, 2) \cap (\mathbb{R} \setminus \mathbb{Q}). \end{cases}$$

Which one of the following is not true?

- A)  $f$  is continuous at  $x = 1$
- B)  $f$  is differentiable at  $x = 1$
- C)  $f$  is not differentiable at  $x = 1$
- D)  $f$  is differentiable only at  $x = 1$ .

(31) Let  $R$  be a finite commutative ring with unity and  $P$  be an ideal in  $R$  satisfying:  
 $ab \in P \implies a \in P$  or  $b \in P$ , for any  $a, b \in R$ . Consider the statements:

- (i)  $P$  is a finite ideal
- (ii)  $P$  is a prime ideal
- (iii)  $P$  is a maximal ideal.

Then

- A) (i),(ii) and (iii) are all correct
- B) None of (i),(ii) or (iii) is correct
- C) (i) and (ii) are correct but (iii) is not correct
- D) (i) and (ii) are not correct but (iii) is correct.

(32) Let  $\phi : R \rightarrow R'$  be a non-zero mapping such that  $\phi(a + b) = \phi(a) + \phi(b)$  and  $\phi(ab) = \phi(a)\phi(b)$  for all  $a, b \in R$ , where  $R, R'$  are rings with unity. Then

- A)  $\phi(1) = 1$  for all rings with unity  $R, R'$
- B)  $\phi(1) \neq 1$  for any rings with unity  $R, R'$
- C)  $\phi(1) \neq 1$  if  $R'$  is an integral domain or if  $\phi$  is onto
- D)  $\phi(1) = 1$  if  $R'$  is an integral domain or if  $\phi$  is onto.

(33) Let  $R$  be a ring,  $L$  be a left ideal of  $R$  and let  $\lambda(L) = \{x \in R \mid xa = 0 \forall a \in L\}$ .

Then

- A)  $\lambda(L)$  is not a two-sided ideal of  $R$
- B)  $\lambda(L)$  is a two-sided ideal of  $R$
- C)  $\lambda(L)$  is a left but not right ideal of  $R$
- D)  $\lambda(L)$  is a right but not left ideal of  $R$ .

(34) Let  $S = \{a + ib \mid a, b \in \mathbb{Z}, b \text{ is even}\}$ . Then

- A)  $S$  is both a subring and an ideal of  $\mathbb{Z}[i]$
- B)  $S$  is neither an ideal nor a subring of  $\mathbb{Z}[i]$
- C)  $S$  is an ideal of  $\mathbb{Z}[i]$  but not a subring of  $\mathbb{Z}[i]$
- D)  $S$  is a subring of  $\mathbb{Z}[i]$  but not an ideal of  $\mathbb{Z}[i]$ .

- (35) The set of all ring homomorphism  $\phi : \mathbb{Z} \rightarrow \mathbb{Z}$
- A) is an infinite set
  - B) has exactly two elements
  - C) is a singleton set
  - D) is an empty set.
- (36) Let  $F$  be a field of characteristic 2. Then
- A) either  $F$  has  $2^n$  elements or is an infinite field
  - B)  $F$  is an infinite field
  - C)  $F$  is a finite field with  $2^n$  elements
  - D) either  $F$  is an infinite field or a finite field with  $2n$  elements.
- (37) Consider the following classes of commutative rings with unity: ED is the class of Euclidean domain, PID is the class of principal ideal domain, UFD is the class of unique factorization domain and ID is the class of integral domain. Then
- A)  $\text{PID} \subset \text{ED} \subset \text{UFD} \subset \text{ID}$
  - B)  $\text{ED} \subset \text{UFD} \subset \text{PID} \subset \text{ID}$
  - C)  $\text{ED} \subset \text{PID} \subset \text{UFD} \subset \text{ID}$
  - D)  $\text{UFD} \subset \text{PID} \subset \text{ED} \subset \text{ID}$ .
- (38) Consider the polynomial ring  $\mathbb{Z}[x]$  and  $\mathbb{Q}[x]$ . Then
- A)  $\mathbb{Z}[x]$  and  $\mathbb{Q}[x]$  both are Euclidean domains
  - B)  $\mathbb{Z}[x]$  and  $\mathbb{Q}[x]$  both are not Euclidean domains
  - C)  $\mathbb{Z}[x]$  is a Euclidean domain but  $\mathbb{Q}[x]$  is not a Euclidean domain
  - D)  $\mathbb{Q}[x]$  is a Euclidean domain but  $\mathbb{Z}[x]$  is not a Euclidean domain.
- (39) Let  $R$  be a commutative ring with unity such that the polynomial ring  $R[x]$  is a principal ideal domain. Then
- A)  $R$  is a field
  - B)  $R$  is a PID but not a field
  - C)  $R$  is a UFD but not a field
  - D)  $R$  is not a field but is an integral domain.
- (40) Let  $T$  be a linear transformation on  $\mathbb{R}^3$  defined by  $T(x_1, x_2, x_3) = (3x_1, x_1 - x_2, 2x_1 + x_2 + x_3)$ . What is  $T^{-1}$ ?
- A)  $T^{-1}(x_1, x_2, x_3) = \left(\frac{x_1}{3}, \frac{x_1}{3} + x_2, -x_1 + x_2 + x_3\right)$
  - B)  $T^{-1}(x_1, x_2, x_3) = \left(\frac{x_1}{3}, \frac{x_1}{3} - x_2, x_1 + x_2 + x_3\right)$
  - C)  $T^{-1}(x_1, x_2, x_3) = \left(\frac{x_1}{3}, \frac{x_1}{3} - x_2, -x_1 + x_2 + x_3\right)$
  - D)  $T^{-1}(x_1, x_2, x_3) = \left(\frac{x_1}{3}, \frac{x_1}{3} + x_2, x_1 + x_2 + x_3\right)$ .

- (41) Let  $V$  be the vector space of all  $n \times n$  matrices over a field  $F$ . Which one of the following is not a subspace of  $V$ ?
- A) All upper triangular matrices of order  $n$   
 B) All non-singular matrices of order  $n$   
 C) All symmetric matrices of order  $n$   
 D) All matrices of order  $n$ , the sum of whose diagonal entries is zero.
- (42) Let  $V$  be the vector space of all  $n \times n$  matrices over a field. Let  $V_1$  be the subspace of  $V$  consisting of all symmetric matrices of order  $n$  and  $V_2$  be the subspace of  $V$  consisting of all skew-symmetric matrices of order  $n$ . Which one of the following is not a subspace of  $V$ ?
- A)  $V_1 + V_2$       B)  $V_1 \cup V_2$       C)  $V_1 \oplus V_2$       D)  $V_1 \cap V_2$ .
- (43) Let  $V = \mathbb{R}^3$  be the real inner product space with the usual inner product. A basis for the subspace  $u^\perp$  of  $V$ , where  $u = (1, 3, -4)$ , is
- A)  $\{(1, 0, 3), (0, 1, 4)\}$ ,      B)  $\{(3, -1, 0), (-6, 2, 0)\}$   
 C)  $\{(-3, 1, 0), (4, 0, 1)\}$       D)  $\{(3, 1, 0), (-4, 0, 1)\}$ .
- (44) The matrix  $A$  that represents the linear operator  $T$  on  $\mathbb{R}^2$ , where  $T$  is the reflection in  $\mathbb{R}^2$  about the line  $y = -x$  is
- A)  $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$   
 B)  $A = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$   
 C)  $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$   
 D)  $A = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$ .
- (45) Consider the subspace  $U$  of  $\mathbb{R}^4$  spanned by the vectors  $v_1 = (1, 1, 1, 1)$ ,  $v_2 = (1, 1, 2, 4)$ ,  $v_3 = (1, 2, -4, -3)$ . An orthonormal basis of  $U$  is
- A)  $\{\frac{1}{2}(1, 1, 1, 1), \frac{1}{\sqrt{6}}(-1, -1, 0, 2), \frac{1}{\sqrt{2}}(1, 3, -6, 2)\}$   
 B)  $\{\frac{1}{2}(1, 1, 1, 1), \frac{1}{2\sqrt{6}}(-1, -1, 0, 2), \frac{1}{\sqrt{2}}(1, 3, 6, -2)\}$   
 C)  $\{\frac{1}{2}(1, 1, 1, 1), \frac{1}{\sqrt{6}}(-1, -1, 0, 2), \frac{1}{5\sqrt{2}}(1, 3, -6, 2)\}$   
 D)  $\{(1, 1, 1, 1), (-1, -1, 0, 2), (1, 3, -6, 2)\}$ .
- (46) Let  $V$  be a vector space over  $\mathbb{Z}_5$  of dimension 3. The number of elements in  $V$  is
- A) 5      B) 125      C) 243      D) 3.



- (47) Let  $W$  be the subspace of  $\mathbb{R}^4$  spanned by the vectors  $u_1 = (1, -2, 5, -3)$ ,  $u_2 = (2, 3, 1, -4)$ ,  $u_3 = (3, 8, -3, -5)$ . The dimension of  $W$  is
- A) 1                      B) 2                      C) 3                      D) 4.
- (48) Let  $\lambda$  be a non-zero characteristic root of a non-singular matrix  $A$  of order  $2 \times 2$ . Then a characteristic root of the matrix  $\text{adj.}A$  is
- A)  $\frac{\lambda}{|A|}$                       B)  $\frac{|A|}{\lambda}$                       C)  $\lambda|A|$                       D)  $\frac{1}{\lambda}$ .
- (49) Let  $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  be a  $2 \times 2$  matrix. Then the expression  $A^5 - 2A^4 - 3A^3 + A^2$  is equal to
- A)  $2A + 3I$                       B)  $3A + 2I$                       C)  $2A - 3I$                       D)  $3A - 2I$ .
- (50) The number of elements in the group  $\text{Aut } \mathbb{Z}_{200}$  of all automorphisms of  $\mathbb{Z}_{200}$  is
- A) 78                      B) 80                      C) 84                      D) 82.
- (51) Let  $A = \begin{pmatrix} 2 & 6 \\ 3 & 5 \end{pmatrix}$  be a matrix over the integers modulo 11. The inverse of  $A$  is
- A)  $A = \begin{pmatrix} 8 & 9 \\ 10 & 9 \end{pmatrix}$   
 B)  $A = \begin{pmatrix} 10 & 8 \\ 9 & 9 \end{pmatrix}$   
 C)  $A = \begin{pmatrix} 9 & 10 \\ 9 & 8 \end{pmatrix}$   
 D)  $A = \begin{pmatrix} 9 & 9 \\ 10 & 8 \end{pmatrix}$ .
- (52) The order of the group  $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid ad - bc = 1 \text{ and } a, b, c, d \in \mathbb{Z}_3 \right\}$  relative to matrix multiplication is
- A) 18                      B) 20                      C) 24                      D) 22.
- (53) The number of subgroups of the group  $\mathbb{Z}_{200}$  is
- A) 8                                      B) 14  
 C) 12                                      D) 10.

(54) Let  $G = U(32)$  and  $H = \{1, 31\}$ . The quotient group  $G/H$  is isomorphic to

- A)  $\mathbb{Z}_8$
- B)  $\mathbb{Z}_4 \oplus \mathbb{Z}_2$
- C)  $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2$
- D) The dihedral group  $D_4$ .

(55) The number of sylow 5-subgroups of the group  $\mathbb{Z}_6 \oplus \mathbb{Z}_5$  is

- A) 6
- B) 4
- C) 12
- D) 1.

(56) The singular solution of the first order differential equation  $p^3 - 4xyp + 8y^2 = 0$  is

- A)  $27x - 4y^3 = 0$
- B)  $27y - 4x^2 = 0$
- C)  $27y - 4x^3 = 0$
- D)  $27y + 4x^3 = 0$ .

(57) The general solution of the system of first order differential equations

$$\frac{dx}{dt} + \frac{dy}{dt} = x + t,$$

$$\frac{dx}{dt} - \frac{d^2y}{dt^2} = 0$$

is given by

- A)  $x = \frac{1}{2}t + c_1t^2 + c_2t$ ;  $y = \frac{1}{2}t - c_1t + c_2$
- B)  $x = \frac{1}{2}t^2 + c_1t + c_2$ ;  $y = \frac{1}{6}t^3 + \frac{1}{2}c_1t^2 + (c_2 - c_1)t + c_3$
- C)  $x = \frac{1}{2}t^2 - c_1t + c_2t^2$ ;  $y = \frac{1}{6}t^2 + \frac{1}{2}c_1t^2 + (c_2 - c_1)t^2 + c_3$
- D)  $x = \frac{1}{3}t^2 + c_1t + c_2$ ;  $y = \frac{1}{6}t^3 - \frac{1}{2}c_1t + (c_2 - c_1)t^2 + c_3$ .

(58) Consider the following statements regarding the two solutions  $y_1(x) = \sin x$  and  $y_2(x) = \cos x$  of  $y'' + y = 0$ :

- (i) They are linearly dependent solutions of  $y'' + y = 0$ .
  - (ii) Their wronskian is 1.
  - (iii) They are linearly independent solutions of  $y'' + y = 0$ .
- which of the statements is true?

- A) (i) and (ii)
- B) (ii) and (iii)
- C) (iii)
- D) (i).

(59) The general solution of  $\frac{d^4y}{dx^4} - 5\frac{d^3y}{dx^3} + 6\frac{d^2y}{dx^2} + 4\frac{dy}{dx} - 8y = 0$  is

- A)  $y = c_1 + c_2x + c_3x^2 + c_4e^x$
- B)  $y = c_1 - c_2x + c_3x^3 + c_4e^{-x}$
- C)  $y = (c_1 + c_2x + c_3x^2)e^{2x} + c_4e^x$

D)  $y = (c_1 + c_2x + c_3x^2)e^{2x} + c_4e^{-x}$ .

(60) The solution of the initial value problem  $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 25y = 0$ ,  $y(0) = -3$ ,  $y'(0) = -1$  is

A)  $y = e^{3x}(2 \cos 4x + 3 \sin 2x)$

B)  $y = e^{-3x}(2 \sin 2x - 3 \cos 2x)$

C)  $y = e^{3x}(2 \sin 4x - 3 \cos 4x)$

D)  $y = e^{3x}(2 \sin 4x + 3 \cos 4x)$ .

(61) The Sturm-Liouville problem given by  $y'' + \lambda y = 0$ ,  $y(0) = 0$ ,  $y(\pi) = 0$  has a trivial solution if

A)  $\lambda \leq 0$

B)  $\lambda > 0$

C)  $0 < \lambda < 1$

D)  $\lambda \geq 1$ .

(62) The initial value problem  $y' = 1 + y^2$ ,  $y(0) = 1$  has the solution given by

A)  $y = \tan(x - \frac{\pi}{4})$

B)  $y = \tan(x + \frac{\pi}{4})$

C)  $y = \tan(x - \frac{\pi}{2})$

D)  $y = \tan(x + \frac{\pi}{2})$ .

(63) The series expansion that gives  $y$  as a function of  $x$  in neighborhood of  $x = 0$  when  $\frac{dy}{dx} = x^2 + y^2$ ; with boundary conditions  $y(0) = 0$  is given by

A)  $y = \frac{1}{3}x^3 + \frac{1}{63}x^7 + \frac{2}{2079}x^{11} + \dots$

B)  $y = \frac{1}{2}x^3 + \frac{1}{8}x^5 + \frac{1}{32}x^7 + \dots$

C)  $y = x^2 + \frac{1}{2!}x^3 + \frac{1}{3!}x^4 + \dots$

D)  $y = \frac{1}{2!}x^2 + \frac{1}{4!}x^4 + \frac{1}{6!}x^6 + \dots$ .

(64) The value of  $y(0.2)$  obtained by solving the equation  $\frac{dy}{dx} = \log(x + y)$ ,  $y(0) = 1$  by modified Euler's method is equal to

A) 1.223

B) 1.0082

C) 2.381

D) 1.639.

(65) Reciprocal square root iteration formula for  $N^{-1/2}$  is given by

A)  $x_{i+1} = \frac{x_i}{2}(3 - x_i^2N)$

B)  $x_{i+1} = \frac{x_i}{9}(4 - x_i^2N)$

C)  $x_{i+1} = \frac{1}{16}(8 - x_i^2N)$

D)  $x_{i+1} = \frac{x_i}{4}(10 - x_i^2N)$ .



- A)  $z(x, t) = \sum_{n=0}^{\infty} c_n \cos(nx + \epsilon_n) e^{-n^2 kt}$   
 B)  $z(x, t) = \sum_{n=0}^{\infty} c_n \cos(nx + \epsilon_n) e^{n^2 kt}$   
 C)  $z(x, t) = \sum_{n=0}^{\infty} c_n \sin(nx + \epsilon_n) e^{n^2 kt}$   
 D)  $z(x, t) = \sum_{n=-\infty}^{\infty} c_n \sin(nx + \epsilon_n) e^{n^2 kt}$ .

(72) The complete integral of the equation  $p^2 y(1 + x^2) = qx^2$  is

- A)  $z = a(1 + x^2) + \frac{1}{2}a^2 y^2 + b$   
 B)  $z = \frac{1}{2}a^2 \sqrt{1 + x^2} + a^2 y^2 + b$   
 C)  $z = a\sqrt{1 + x^2} + \frac{1}{2}a^2 y^2 + b$   
 D)  $z = a(1 + x^2) + \frac{1}{2}ay + b$ .

(73) The general integral of the partial differential equation  $z(xp - yq) = y^2 - x^2$  is

- A)  $x^2 + y^2 + z^2 = f(xy)$   
 B)  $x^2 - y^2 + z^2 = f(xy)$   
 C)  $x^2 - y^2 - z^2 = f(xy)$   
 D)  $x^2 + y^2 - z^2 = f(xy)$ .

(74) The solution of the partial differential equation  $\frac{\partial^4 z}{\partial x^4} + \frac{\partial^4 z}{\partial y^4} = 2 \frac{\partial^4 z}{\partial x^2 \partial y^2}$  is

- A)  $z = x\phi_1(x + y) + \phi_2(x + y) + x\psi_1(x + y) + \psi_2(x + y)$   
 B)  $z = x\phi_1(x - y) + \phi_2(x - y) + x\psi_1(x - y) + \psi_2(x - y)$   
 C)  $z = x\phi_1(x + y) + \phi_2(x - y) + x\psi_1(x + y) + \psi_2(x - y)$   
 D)  $z = x\phi_1(x - y) + \phi_2(x - y) + x\psi_1(x + y) + \psi_2(x + y)$ .

(75) The eigen values and eigen functions of the vibrating string problem  $u_{tt} - c^2 u_{xx} = 0$ ,  $0 \leq x \leq l$ ,  $t > 0$ ,  $u(x, 0) = f(x)$ ,  $0 \leq x \leq l$ ,  $u_t(x, 0) = g(x)$ ,  $0 \leq x \leq l$ ,  $u(0, t) = 0$ ,  $u(l, t) = 0$ ,  $t \geq 0$  are

- A)  $(\frac{n\pi}{l})^2, \sin \frac{n\pi x}{l}, n = 1, 2, 3, \dots$   
 B)  $(\frac{n\pi}{l})^2, \cos \frac{n\pi x}{l}, n = 1, 2, 3, \dots$   
 C)  $\frac{n\pi}{l}, \sin \frac{n\pi x}{l}, \cos \frac{n\pi x}{l}, n = 1, 2, 3, \dots$   
 D) All the above.